



The biodegradability of a feedstock is determining the optimal C/N ratios in anaerobic digestion

Kaldis, Fokion ; Karakashev, Dimitar; Angelidaki, Irini

Published in:
Book of Abstracts. DTU's Sustain Conference 2015

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Kaldis, F., Karakashev, D., & Angelidaki, I. (2015). The biodegradability of a feedstock is determining the optimal C/N ratios in anaerobic digestion. In *Book of Abstracts. DTU's Sustain Conference 2015* [R-18] Technical University of Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The biodegradability of a feedstock is determining the optimal C/N ratios in anaerobic digestion.

Fokion Kaldis^{1,2}, Dimitar Karakashev², Irini Angelidaki^{2*} (1) Biological Applications and Technologies, University of Ioannina, Ioannina, Greece, (2) Department of Environmental Engineering, Technical University of Denmark

*Email: iria@env.dtu.dk

Anaerobic digestion is a collection of processes by which different types of microorganisms break down biodegradable material in the absence of oxygen. Microorganisms need nitrogen for building up their cell mass. Approx. 15% of the cell mass is nitrogen. Therefore nitrogen is a necessary nutrient for growth of microorganisms. On the other hand excessive amounts of nitrogen in the form of ammonia are toxic for the microbial growth (Xiaojiang et al 2012 and Kayhanian, 1999). In the past main focus has been paid on the carbon to nitrogen (C/N) ratio as an important parameter for design and operation of AD process. However, this conception is a simplification of the reality and is not connected with any mechanistic explanation. Biomass is degraded to build new cells, with a cell to substrate yield depending on the microorganism and the energy content of the feedstock biomass used as substrate. For a successful balanced process in respect to C/N ratio, we would need exact balance of nitrogen to adequately cover the nutritional needs of the microorganisms, but not excessive to create inhibition problems. Therefore, we hypothesise that the biodegradability of the feedstock is a very important factor which has been ignored so far, in the optimisation of the C/N ratio of the feedstocks.

In order to test our hypothesis we have investigated feedstocks with different biodegradability in order to determine the effect of biodegradability on the C/N ratio, and to establish the connection of the C/N ratio with the cell biomass composition. As model-substrates (feedstock biomasses) with different biodegradability we tested: glucose (easy degradable) and grass (rich in lignocellulose and difficult to biodegrade). Based on literature data for optimal, sub-optimal and super-optimal ranges for C/N ratio following ratio were chosen: 5, 20, and 35 and 50. All experiments took place under thermophilic (55°C) conditions due to its superiority when compared to the mesophilic conditions for the bio-conversion of the substrates into bio-methane. Results obtained can be utilized for development of innovative co-digestion process configuration with substrates with different biodegradabilities

References:

Xiaojiang W., Gaihe Y., Feng Y., Guangxin R., Xinhui H., 2012. Optimizing feeding composition and carbon-nitrogen ratios for improved methane yield during anaerobic co-digestion of dairy, chicken manure and wheat straw. *Bioresource Technology* ; 120:78-83.

Kayhanian M., 1999. Ammonia inhibition in high-solids biogasification: an overview and practical solutions *Environ. Technol.*, 20, pp. 355–365